



DR. WISAM MENESI, MD, FRCSC

Cosmetic and Reconstructive Plastic Surgery

**103-457 Bishop Drive
Fredericton, NB
E3C 2M6**

**Telephone: (506) 455-1188
Fax: (506) 455-1189**

Postoperative Instructions for Blepharoplasty

INSTRUCTIONS

- Have someone drive you home after surgery and help you at home for 1-2 days.
- Follow balanced diet.
- Decreased activity may promote constipation, add fruit to your diet and increase fluid intake.
- Take pain medication as prescribed. Do not take aspirin or any products containing aspirin.
- Do not drink alcohol when taking pain medications.
- No alcohol for 3 weeks.
- Do not smoke, as smoking delays healing and increases the risk of complications.

ACTIVITIES

- Start walking as soon as possible, as this helps to reduce swelling and lowers the chance of blood clots.
- Do not drive until you are no longer taking any pain medications (narcotics).
- Avoid vigorous exercise and activities are restricted for at least 3 weeks. No heavy lifting, pulling or bending.
- Return to work in 2 weeks (Depends on job).

INCISION CARE

- Keep surgical tapes in place until they start to fall off themselves.
- Keep incisions dry for 24 hours, then you can wash your face gently with soap and water, then pat it dry.
- Keep your head elevated for several days and sleep with your head on 2 pillows for at least 1 week.
- Avoid rubbing your eyes.
- Avoid exposing scars to sun for at least 12 months.
- Always use sunblock, if sun exposure is unavoidable (SPF 30 or greater).

WHAT TO EXPECT

- You may experience temporary sensitivity around the incisions for few days.
- wear sunglasses when outdoor to protect your eyes.
- You may have bruising and swelling. The majority of bruising and swelling will subside in 1 week.
- Scars will be reddened for few weeks. They will fade and soften over time, it takes up to 6-12 months for the scars to mature.

FOLLOW-UP CARE

- Sutures will usually be removed in 1 week.

WHEN TO CALL

- If you have sudden painful swelling or bruising.
- If swelling and redness persist after a few days.
- If you have increased redness along the incision.
- If you have severe or increased pain not relieved by medication.
- If you have any yellowish or greenish drainage from the incisions or notice a foul odor.
- If you have bleeding from the incisions that is difficult to control with light pressure.